

SURFACES WITH $p_g = q = 1$, $K^2 = 8$ AND NONBIRATIONAL BICANONICAL MAP

GIUSEPPE BORRELLI

ABSTRACT. We prove that if the bicanonical map of a minimal surface of general type S with $p_g = q = 1$ and $K_S^2 = 8$ is non birational, then it is a double cover onto a rational surface.

An application of this theorem is the complete classification of minimal surfaces of general type with $p_g = q = 1$, $K_S^2 = 8$ and nonbirational bicanonical map.

Let S be a smooth surface and K_S a canonical divisor of S . The m -th canonical map is the rational map φ_{mK_S} associated to the complete linear system $|mK_S|$. Investigating the behavior of φ_{mK_S} ($m > 0$) is a useful tool when studying surfaces of general type, i.e. surfaces for which φ_{mK_S} is generically finite for some $m \geq 1$.

The behavior of the maps φ_{mK_S} is well understood when $m \geq 3$ (cf. [11] for a survey). For instance, we know that if $m \geq 5$, then φ_{2K_S} maps S birationally onto a normal surface (cf. [5]).

Let S be a (smooth) minimal surface of general type. Assume that the bicanonical (i.e. $m = 2$) map is non birational. Then S may or may not admit a fibration whose general fiber is a smooth curve of genus 2. If it does, we say that S *presents the standard case*. By [19] we know that if $K_S^2 > 9$, then S presents the standard case.

Surfaces with non birational bicanonical map not presenting the standard case have been intensively investigated in the last decade by several authors, and their classification is rather understood with the exception of the cases when $p_g = q \leq 1$ (cf. [10] for a survey on this subject).

The cases when $p_g = q = 0$ have been studied by Mendes Lopes and Pardini in the last years. We refer to [13] and [14] for a survey, here we just mention that they obtained a complete classification of surfaces with $K_S^2 \geq 6$ and nonbirational bicanonical map. It follows from their analysis that if $K_S^2 \geq 6$ and φ_{2K} is nonbirational, then $K_S^2 \leq 8$ and φ_{2K} factors through a double cover onto a rational surface. Furthermore, $\deg \varphi_{2K} = 2$ if $K_S^2 = 7, 8$ and $\deg \varphi_{2K} = 2$ or 4 if $K_S^2 = 6$.

Let S be a minimal surface of general type with $p_g = q = 1$. Then $K_S^2 \geq 2$ by [5, Lemma 14] and $K_S^2 \leq 9$ by [15] and [24]. The only cases for which there is a complete classification from the point of view of the non birationality of the bicanonical map are those when $K_S^2 = 2$ and $K_S^2 = 9$. In the first case S presents the standard case, in fact the Albanese map is a fibration with general fiber a curve of genus 2 (cf. [8]). If the second case occurs, then the bicanonical map is birational (cf. [9, Proposition 3.1] and [23, Théorème 2.2 and Corollaire 3]).

Surfaces with $K_S^2 = 8$ have been studied by Polizzi in [18], who considered those whose bicanonical map factors through a double cover onto rational surface. He proved that in this case φ_{2K} has degree 2 and that these surfaces are quotient of the

product of two curves by the action of a finite group. He also studied the moduli space and showed that these surfaces in fact exist.

In [18] it is also claimed that if the bicanonical map of a minimal surface with $p_g = q = 1$ and $K_S^2 = 8$ factors through a double cover onto a surface X , then X is rational. For this assertion the author refers to a theorem by Xiao ([22, Theorem 3]), which unfortunately has a gap, as pointed out by Rito (cf. [20]).

In the present paper we prove the following.

Theorem 0.1. *Let S be a minimal surface of general type with $p_g = q = 1$ and $K_S^2 = 8$. If the bicanonical map of S is nonbirational then φ_{2K} is a double cover onto a rational surface.*

An application of this theorem is the complete classification of surfaces with $p_g = q = 1$, $K_S^2 = 8$ and nonbirational bicanonical map (see Theorem 3.1).

The paper is organized as follows. In Section 1 we recall some facts which will be used in the paper. In Section 2 we first fix the gap in [18], and then we study the case when the bicanonical map has degree 4. In the last section, first we prove Theorem 0.1 and then, as application, we provide the classification of surfaces with $p_g = q = 1$, $K_S^2 = 8$ and nonbirational bicanonical map.

0.1. Notation. Given a smooth surface X , we denote by K_X a canonical divisor of X and use the standard notation $p_g(X) = h^0(K_X)$, $q(X) = h^1(K_X)$ and $\chi(X) = p_g(X) - q(X) + 1$. Throughout the paper S is a minimal surface of general type. For brevity sake, we let $p_g = p_g(S)$ and $q = q(S)$. We use the symbol $\equiv$ (resp. $\sim$) to denote linear (numeric) equivalence of divisors. As usual, a $(-n)$ -curve on a surface is a smooth rational curve with self intersection $-n$.

A *fibration* is a surjective morphism from a smooth surface onto a smooth curve with connected fibers. A fibration is *isotrivial* if the smooth fibers are all isomorphic. A fibration is said *relatively minimal* if the fibers contain no (-1) -curves.

Given a surface X and an effective divisor $D \in \text{Pic}(X)$, we denote by $\varphi_D : X \dashrightarrow \mathbb{P}^{h^0(D)-1}$ the rational map defined by the complete linear system $|D|$.

We will use freely the facts concerning involutions on surfaces and on double covers between surfaces. We refer to [6, section 1.1], from where we keep some definition (for instance, of $[r, r]$ -points in the proof of Proposition 2.1).

1. USEFUL FACTS

We gather some facts which will be used throughout.

Lemma 1.1. *Let X, Y be smooth surfaces and $X \rightarrow Y$ a double cover. If $p_g(Y) = 1$ and $q(X) - q(Y) \geq 3$, then there exists a fibration $X \rightarrow C$ onto a smooth curve of genus $g(C) \geq 2$.*

Proof. Let σ be the involution induced on X by the double cover, and $G \subseteq \text{Aut}(X)$ the group of automorphisms generated by σ , so that $X/G = Y$. For any p there exists an isomorphism $H^0(\Omega_X^p)^G \cong H^0(\Omega_Y^p)$, where $H^0(\Omega_X^p)^G \subseteq H^0(\Omega_X^p)$ is the subspace of G -invariant p -forms.

Let $V^- \subset H^0(\Omega_X^1)$ be the subspace of G -anti invariant 1-forms, so that $H^0(\Omega_X^1) = V^- \oplus H^0(\Omega_X^1)^G$. Then $\dim V^- = q(X) - \dim H^0(\Omega_X^1)^G = q(X) - q(Y) \geq 3$ and,

since $p_g(Y) = 1$, the map $\bigwedge^2 V^- \rightarrow H^0(\Omega_X^2)^G \cong H^0(\Omega_Y^2)$ is not injective. Therefore, by the Castelnuovo-de Franchis' Theorem, there is a fibration $X \rightarrow C$ where C is a curve of genus $g(C) \geq 2$. $\square$

We will use the following results, usually with no further reference.

Proposition 1.2 (cf. [3] or [4]). *Let X be a smooth surface and $f : X \rightarrow C$ a relatively minimal fibration onto a curve of genus b . Let F be a general fiber of f and g its genus. Then*

- (1) $K_X^2 \geq 8(g-1)(b-1)$;
- (2) $12\chi(X) - K_X^2 \geq 4(g-1)(b-1)$;
- (3) $q(X) \leq g + b$.

Furthermore, if equality holds in (1) then the fibers of f have constant modulus, if equality holds in (2) every fiber of f is smooth, and if equality holds in (3) then $X \cong F \times C$.

Proposition 1.3 (cf. [21]). *Let $f : X \rightarrow C$ be a relatively minimal isotrivial fibration, with general fiber a curve F of genus $g \geq 1$. Then there exist a curve A and a finite group G acting faithfully and holomorphically on A and F , such that $C \cong A/G$, and there is a birational map $\pi : X \rightarrow (A \times F)/G$ (where G acts diagonally on $A \times F$) for which the following diagram commutes*

$$\begin{array}{ccc} X & \xrightarrow{\pi} & (A \times F)/G \\ \downarrow f & & \downarrow \\ B & \xrightarrow{\cong} & A/G \end{array}$$

Furthermore,

- (1) $q(X) = g(A/G) + g(F/G)$;
- (2) *the natural map $(A \times F)/G \rightarrow F/G$ give rise to an isotrivial fibrations and $f_1 : X \rightarrow F/G$ whose general fiber is isomorphic to A ;*
- (3) *if the only singular fibers of f are multiples of smooth curves, then π is an isomorphism and G operates on $A \times F$ without fixed points (i.e. $(A \times F) \rightarrow X$ is étale).*

Lemma 1.4. *Let S be minimal surface of general type with $p_g = q = 1$ and $K_S^2 = 8$. Let $S \rightarrow C$ be a fibration. Let b the genus of C and g the genus of a general fiber.*

Then $g \geq 3$ ([23, Théorème 2.2 and Corollaire 3]) and $b \leq 1$ (cf. for instance [4]). In particular, if $b = 1$ then C is the Albanese variety of S and $S \rightarrow C$ is the Albanese morphism.

2. THE BICANONICAL MAP OF SURFACES WITH $K_S^2 = 8$

Throughout this section let S be a minimal surface of general type with $p_g = q = 1$, $K_S^2 = 8$ and nonbirational bicanonical map. Let $\varphi_2 = \varphi_{2K_S} : S \dashrightarrow S_2 \subset \mathbb{P}^8$ be the bicanonical map of S . Then φ_2 is a morphism (cf. [19]) and, since by [16] K_S is ample, it is finite. Let $d \geq 2$ be the degree of φ_2 and $\deg(S_2)$ the degree of S_2 . Then $(2K_S)^2 = d \cdot \deg(S_2)$ and so $d = 2$ or 4 , because an irreducible non degenerate surface in $\mathbb{P}^n$ has degree at least $n - 1$.

2.1. The case when φ_2 factors through an involution. The following proposition fix the gap of [18] (cf. the introduction).

Proposition 2.1. *Let S be a minimal surface of general type with $p_g = q = 1$ and $K_S^2 = 8$. Suppose that the bicanonical map of S factors through an involution σ . Then the quotient S/σ is birational to a rational surface.*

Proof. Let $\hat{\Sigma} \rightarrow S/\sigma$ be the minimal desingularization, and suppose that $\hat{\Sigma}$ is not rational. Then, by [6, Theorem 2.7], $\hat{\Sigma}$ has non negative Kodaira dimension and thus, since S does not present the standard case, it is birational to a $K3$ surface (cf. [20]).

Let $\pi : \hat{S} \rightarrow S$ the blow up of the isolated fixed points of σ , and $\hat{\sigma}$ the involution induced on $\hat{S}$. Then the quotient $\hat{S}/\hat{\sigma}$ is isomorphic to $\hat{\Sigma}$ and we have the following commutative diagram

$$\begin{array}{ccc} \hat{S} & \xrightarrow{\pi} & S \\ \rho \downarrow & & \downarrow \\ \hat{\Sigma} & \longrightarrow & S/\sigma \end{array}$$

where ρ is the canonical projection. Let $E_1, \dots, E_\nu$ be the exceptional (-1) -curves of π , and $C_i = \rho(E_i)$. Let $\hat{B}$ be the branch curve of ρ . Then C_i is a (-2) -curve belonging to $\hat{B}$ for every i .

Let $\phi : \hat{\Sigma} \rightarrow \Sigma$ be the birational morphism to the minimal model, and $B = \phi_* \hat{B}$. Then $B \equiv 2\Delta$ for some $\Delta \in \text{Pic}(\Sigma)$ and, by [20, proof of Theorem 1], B has at most one essential singularity, which is either a $[4]$ -point or a $[3, 3]$ -point.

Let $\mathcal{E}_1, \dots, \mathcal{E}_d$ be the reduced and irreducible exceptional curves of ϕ . Since Σ is a $K3$ surface, $K_{\hat{\Sigma}} = \sum e_j \mathcal{E}_j$, where the e_j 's are positive integers. Since the C_i 's are (-2) -curves, we have $C_i \cdot \sum e_j \mathcal{E}_j = 0$ for every i . Hence either $C_i \cap \sum \mathcal{E}_j = \emptyset$ or $C_i = \mathcal{E}_j$ for some j , i.e. either $\phi(C_i)$ is a (-2) -curve or $\phi(C_i)$ is a point. Thus, by [6, Lemma 1.4], there are at least $\nu - 1$ pairwise disjoint (-2) -curves on Σ and then $\nu \leq 17$, because Σ is a $K3$ surface.

By assumption, φ_2 factors through σ . Then, by [6, Proposition 1.8], we get $\nu = K_S^2 - 2\chi(S) + 6\chi(\Sigma) = 18$, which is a contradiction. Therefore S/σ is birational to a rational surface. $\square$

2.2. The case when $d = 4$. In this section we prove the following

Theorem 2.2. *Let S be a minimal surface of general type with $p_g = q = 1$ and $K_S^2 = 8$. Suppose that the degree of the bicanonical map is 4. Then φ_2 factors through a generically finite map of degree two onto a rational surface.*

Assume $d = 4$. Then S_2 is a surface of degree 8 in $\mathbb{P}^8$, and thus it is one of the following (cf. [17]).

- A cone over an elliptic curve of degree 8 in $\mathbb{P}^7$.
- The Veronese embedding in $\mathbb{P}^8$ of a quadric Q in $\mathbb{P}^3$.
- A del Pezzo surface of degree 8, i.e. the image of $\mathbb{P}^2$ by the rational map associated to the linear system $|\mathcal{O}_{\mathbb{P}^2}(3) \otimes \mathcal{I}_x|$, where $\mathcal{I}_x$ is the ideal sheaf of a point in $\mathbb{P}^2$.

We will consider each case separately. In particular, we are going to show that the first case does not occur and that in both the others φ_2 factors through a double cover onto a rational surface.

Theorem 2.2 follows from Lemma 2.3, Propositions 2.4 and 2.6 below.

Lemma 2.3. *S_2 is not a cone over an elliptic curve.*

Proof. Suppose that S_2 is a cone over an elliptic curve C . Let $\mu : S_2 \dashrightarrow C$ be the projection from the vertex. The composite map $f = \mu \circ \varphi_2 : S \dashrightarrow C$ is surjective and thus a morphism, since C has positive genus.

The Stein factorization of f yields a fibration $\tilde{f} : S \rightarrow \tilde{C}$ over a smooth curve of genus $g(\tilde{C}) \geq 1$, whose general fiber is a curve F of genus $g(F)$. Since $K_S^2 = 8$, by Lemma 1.4 we have $g(F) \geq 3$ and thus $F \cdot K_S \geq 4$.

Let $\ell = \varphi_2(F)$. Then ℓ is a general line on S_2 and, since K_S is ample, we have $\varphi_2^* \ell \sim aF$, where $a \geq 1$ is an integer. But then we get $4 = (2K_S) \cdot (aF) \geq 8a$, which is absurd. $\square$

Proposition 2.4. *Suppose that S_2 is the Veronese embedding in $\mathbb{P}^8$ of a quadric Q in $\mathbb{P}^3$. Then the bicanonical map of S factors through a double cover onto a rational surface.*

Proof. Let H be the pull back to S of a general hyperplane section of Q , so that $2H \equiv 2K_S$. Let $\eta = H - K_S$. Then $2\eta \equiv 0$ and so, since $h^0(K_S + \eta) = h^0(H) = 4 \neq h^0(K_S)$, η is a non zero 2-torsion element.

Let $\psi : X \rightarrow S$ be the étale double cover defined by $2\eta \equiv 0$. We have $\chi(X) = 2\chi(S) = 2$, $K_X^2 = 2K_S^2 = 16$ and $p_g(X) = p_g(S) + h^0(K_S + \eta) = 5$. Whence $q(X) = 4$ and thus, by Proposition 1.1, there exists a fibration $f : X \rightarrow C$ over a curve of genus $g(C) \geq 2$. Let F be a general fiber of f and $g(F)$ its genus. Since X is of general type, $g(F) \geq 2$. Observe that, since $\psi : X \rightarrow S$ is étale and S is minimal, X is minimal. In particular, the fibration $f : X \rightarrow C$ is relatively minimal.

Claim. *The fibers of $f : X \rightarrow C$ are smooth and isomorphic curves of genus 3, and we have $g(C) = 2$.*

Suppose $g(F) = 2$. Since $q(X) = 5$, by Proposition 1.2 we get $g(C) \geq 3$. Let σ be the involution on X induced by the double cover $\psi : X \rightarrow S$. Since F does not dominate C , $\sigma(F)$ is a fiber of f . Then there are a fibration $f' : S \rightarrow C'$ over a smooth curve whose general fiber is $\psi(F)$ and a finite morphism $h : C \rightarrow C'$ such that $f' \circ \psi = h \circ f$. Since the genus of $\psi(F)$ is at most $g(F) = 2$, we contradicts Lemma 1.4. Therefore, $g(F) \geq 3$.

By Proposition 1.2, we have $16 = K_X^2 \geq 8(g(F) - 1)(g(C) - 1)$ which yields $g(F) = 3$ and $g(C) = 2$. Since $12\chi(X) - K_X^2 = 8 = 4(g(F) - 1)(g(C) - 1)$, the fibers are all smooth and isomorphic. This proves the claim.

By Proposition 1.3 there are a curve A and a finite group G acting faithfully and holomorphically on F and A , such that $X \cong (A \times F)/G$. Furthermore, there exists a fibration $f_1 : X \rightarrow F/G$ whose general fiber, say $\hat{A}$, is isomorphic to A . In particular, we have $F \cdot \hat{A} = |G|$.

Let $C_1 = F/G$. Since $4 = q(X) = g(A/G) + g(F/G) = 2 + g(C_1)$, we get $g(C_1) = 2$ and the same argument we used for $f : X \rightarrow C$ yields $g(A) = 3$.

We have $p_g(A \times F) = g(F)g(A) = 9$ and $q(A \times F) = g(F) + g(A) = 6$. Thus $\chi(A \times F) = 4$ and so, since $A \times F \rightarrow X$ is étale, $|G| = 2$. Hence $\hat{A} \cdot F = 2$ and

then f_1 (resp. f) induces a double cover $F \rightarrow C_1$ (resp. $A \rightarrow C$). It follows that F (resp. $\hat{A}$) is a hyperelliptic curve (cf. [1, Lemma 5.10]).

Let τ be the involution on X induced by the hyperelliptic involution of F , and let $Y = X/\tau$ be the quotient. Let σ be the involution induced by the double cover $X \rightarrow S$, and let $N \subseteq \text{Aut}(X)$ be the subgroup generated by τ and σ . We have a commutative diagram

$$\begin{array}{ccc}
 & X & \\
 \psi \swarrow & & \searrow \phi \\
 S = X/\sigma & & Y = X/\tau \\
 \tilde{\psi} \searrow & & \swarrow \tilde{\phi} \\
 & X/N &
 \end{array}$$

where ϕ is the canonical projection, and $\tilde{\psi}, \tilde{\phi}$ are double covers. Note that, since the fibers of $f : X \rightarrow C$ are all isomorphic, τ has not isolated fixed point and hence Y is smooth.

By construction, there is a fibration $f' : Y \rightarrow C$ whose general fiber is $\phi(F)$, which is a smooth rational curve. In particular, Y is not birational to S . Note that since $\phi(A)$ dominates C , $\phi(A)$ is not rational.

Let $\tilde{\sigma}$ be the involution induced on Y by the double cover $Y \rightarrow X/N$. Since $\phi(F)$ does not dominate C , we have that $\tilde{\sigma}(\phi(F))$ is a fiber of $f' : Y \rightarrow C$.

Then there is a fibration $\tilde{f} : X/N \rightarrow \tilde{C}$ and a finite morphism $h : C \rightarrow \tilde{C}$ such that $h \circ f' = \tilde{f} \circ \tilde{\phi}$ and $\tilde{\phi}(\phi(F))$ is a fiber of $\tilde{f}$. Let $\Gamma = \tilde{\phi}(\phi(F))$.

The composite map $\tilde{f} \circ \tilde{\psi} = S \rightarrow \tilde{C}$ is a surjective morphism with general fiber the pull back of Γ . Since $\tilde{\psi}$ is a double cover and Γ is rational, $\tilde{\psi}^*\Gamma$ is connected, i.e. $\tilde{f} \circ \tilde{\psi}$ is a fibration. By Lemma 1.4 we have $g(\tilde{C}) \leq 1$.

By the commutativity of the diagram we have $\psi^*\Gamma = \psi(F)$. Since ψ is étale and on S there are no pencils of genus two curves, $\psi(F) \cong F$ is a hyperelliptic curve of genus 3.

By applying the same argument to the fibration $f_1 : X \rightarrow C_1$ we get a fibration $S \rightarrow \tilde{C}_1$ such that $g(\tilde{C}_1) \leq 1$ whose general fiber is a smooth hyperelliptic curve of genus 3 isomorphic to $\hat{A}$.

It is easy to see that these two fibrations are not the same. Therefore, since on S there exists exactly one fibration over an elliptic curve, namely the Albanese fibration, we have $C \cong \mathbb{P}^1$ or $C_1 \cong \mathbb{P}^1$. By [7] the bicanonical map of S factors through a double cover onto a rational surface. $\square$

Finally we consider the last case. To begin with, let us introduce some notation.

From now on we assume that S_2 is the image of $\mathbb{P}^2$ in $\mathbb{P}^8$ by the birational map associated to the linear system $|\mathcal{O}_{\mathbb{P}^2}(3) \otimes \mathcal{I}_x|$, where $\mathcal{I}_x$ is the ideal sheaf of a point $x \in \mathbb{P}^2$. Let $\hat{\mathbb{P}} \rightarrow \mathbb{P}^2$ be the blow up of x , $\ell \subset \hat{\mathbb{P}}$ the pull back of a general line in $\mathbb{P}^2$, and $\hat{\ell}$ the strict transform of a general line through x . Then S_2 is the image

of $\hat{\mathbb{P}}$ by the birational morphism $\mu : \hat{\mathbb{P}} \rightarrow \mathbb{P}^8$ associated to $|2\ell + \hat{\ell}|$. Note that $|2\ell + \hat{\ell}| = |-K_{\hat{\mathbb{P}}}|$.

Let L and $\hat{L}$ be the pull back to S of $\mu_*\ell$ and $\mu_*\hat{\ell}$, respectively, so that $2K_S \in |2L + \hat{L}|$. We have $\hat{L}^2 = 0$ and $\hat{L} \cdot K_S = 4$. Since $\hat{L}$ is a fiber of the composite morphism $\varphi_{\hat{\ell}} \circ \varphi_2 : S \rightarrow \mathbb{P}^1$, it follows from Lemma 1.4 that $\hat{L}$ is a smooth connected curve of genus 3. In particular, $\varphi_{\hat{L}} = \varphi_{\hat{\ell}} \circ \varphi_2 : S \rightarrow \mathbb{P}^1$ is a fibration.

Fix a smooth curve $B \in |\hat{L}|$, and let $\psi : X \rightarrow S$ be the double cover defined by $B \equiv 2(K_S - L)$. Then X is a smooth surface with $\chi(X) = 3$ and $K_S^2 = 24$. Note that, since S is minimal of general type and the branch curve is a smooth curve of genus 3, X is a minimal surface of general type.

In order to compute the invariants of X we observe that $\varphi_2(\ell)$ is a twisted cubic in $\mathbb{P}^3$. Then $h^0(2K_S - L) = h^0(2K_S) - 4 = 5$, and so

$$p_g(X) = h^0(K_S + (K_S - L)) + p_g(S) = 6.$$

Whence $q(X) = 4$ and thus, by Proposition 1.1, there is a fibration $f : X \rightarrow \mathbb{B}$ onto a smooth curve $\mathbb{B}$ of genus $g(\mathbb{B}) \geq 2$. Let F be a general fiber of f and g its genus. By Proposition 1.2 we have the following possibilities

$$(1) \quad \begin{aligned} g(F) &= 4 \text{ and } g(\mathbb{B}) = 2; \\ g(F) &= 3 \text{ and } g(\mathbb{B}) = 2; \\ g(F) &= 2 \text{ and } 2 \leq g(\mathbb{B}) \leq 4. \end{aligned}$$

Consider the composite morphism $\varphi_{\hat{L}} \circ \psi : X \rightarrow \mathbb{P}^1$. One of the following cases occurs.

- (†) $h_1 = \varphi_{\hat{L}} \circ \psi : X \rightarrow \mathbb{P}^1$ is a fibration whose general fiber is $\psi^*\hat{L}$;
- (‡) there exist a fibration $h_2 : X \rightarrow C$ and a double cover $\tilde{\psi} : C \rightarrow \mathbb{P}^1$ such that $h_2 \circ \tilde{\psi} = \varphi_{\hat{L}} \circ \psi$ and a general fiber of h_2 is isomorphic to $\hat{L}$.

Let M be a general fiber of h_i , so that $g(M) = 5$ in the case (†) and $g(M) = 3$ in the case (‡).

We will proceed by analyzing these two cases separately. We will show that the first one does not occur and that in the second case $\hat{L}$ is hyperelliptic. Before going further let us observe that in any case, since

$$\left(\frac{F}{g(F) - 1} + \frac{M}{g(M) - 1} - \frac{K_X}{6} \right) \cdot K_X = 0,$$

by the algebraic index theorem we have

$$(2) \quad 3(F \cdot M) \leq (g(F) - 1)(g(M) - 1),$$

and the equality holds if and only if

$$6(g(M) - 1)F + 6(g(F) - 1)M \sim (g(F) - 1)(g(M) - 1)K_X.$$

Lemma 2.5. *The case (†) does not occur.*

Proof. To rise a contradiction, suppose that h_1 is a fibration. The curve ψ^*B is a double fiber of h_1 . Then $D \cdot M \equiv 0 \pmod{2}$ for any divisor $D \in \text{Pic}(X)$ and in particular, by (1) and (2), we have $F \cdot M \in \{0, 2, 4\}$.

Suppose $F \cdot M = 4$. Since $g(M) = 5$ and $g(F) \leq 4$ (cf. (1)), (2) yields $g(F) = 4$. Then

$$(3) \quad 4F + 3M \sim 2K_X.$$

By (1), we have $g(\mathbb{B}) = 2$ and so, by Proposition 1.2, $f : X \rightarrow \mathbb{B}$ is isotrivial and smooth. Then, there exist a curve A and a group G such that $(F \times A)/G$. Since $q(X) = g(F/G) + g(A/G) = g(F/N) + g(\mathbb{B})$, we have $g(F/G) = 2$.

Let $\mathbb{B}_1 = F/G$ and let $f_1 : X \rightarrow \mathbb{B}_1$ be the natural projection. Then f_1 is an isotrivial fibration whose general fiber, say $\hat{A}$, is isomorphic to A . Note that, since $g(\mathbb{B}_1) = 2$, Proposition 1.2 yields $g(A) \leq 4$. Hence by (3) we get

$$4|G| + 3M \cdot \hat{A} = (4F + 3M) \cdot \hat{A} = 2K_X \cdot \hat{A} = 4(g(A) - 1) \leq 12.$$

Observe that $|G| > 1$, for otherwise $X \cong F \times \mathbb{B}$ and thus $q(X) = g(F) + g(\mathbb{B}) = 6$, which is a contradiction.

It follows that $\hat{A} \cdot M \leq 1$ and then, since $\hat{A} \cdot M \equiv 0 \pmod{2}$, $\hat{A} \cdot M = 0$. Thus M is a fiber of f_1 and so it is isomorphic to A . But this is absurd, because $g(M) = 5$ and $g(A) \leq 4$. Then $F \cdot M \neq 4$.

Since $g(M) = 5$ and $g(F) \leq 4$, $F \cdot M = 0$ cannot occur. Therefore we are left with $F \cdot M = 2$. Since ψ^*B is a double fiber of h_1 , $F \cdot \psi^{-1}(B) = 1$ and so the restriction of $f : X \rightarrow \mathbb{B}$ to B is an isomorphism. Thus $g(\mathbb{B}) = g(B) = g(\hat{L}) = 3$ and, by (1), we get $g(F) = 2$. But this contradicts (2). Therefore $F \cdot M \neq 2$, and so case $(\dagger)$ does not occur. $\square$

Lemma 2.6. *Suppose that S_2 is a del Pezzo surface of degree 8. Then φ_2 factors through a double cover onto a rational surface.*

Proof. By Lemma 2.5 we have a fibration $h_2 : X \rightarrow C$ whose general fiber M is a smooth curve of genus 3 isomorphic to $\hat{L}$, and a double cover $\tilde{\psi} : C \rightarrow \mathbb{P}^1$ such that $\tilde{\psi} \circ h_2 = \varphi_{\hat{L}} \circ \psi$.

By (2) and (1) we have $3(F \cdot M) \leq 2(g(F) - 1) \leq 6$.

Suppose $F \cdot M = 0$. Then M and F are fibers of the same fibration. Hence $g(F) = g(M) = 3$ and, by (1), $C = \mathbb{B}$ is a curve of genus 2. By Riemann-Hurwitz for the double cover $\tilde{\psi} : C \rightarrow \mathbb{P}^1$, the fibration $\varphi_{\hat{L}} : S \rightarrow \mathbb{P}^1$ has at least $(2b + 2) - 1 = 5$ double fibers, say $2\Delta_1, \dots, 2\Delta_5$. The curves $\hat{\ell}_i = \varphi_2(2\Delta_i) \in |\hat{\ell}|$ are reduced, irreducible and distinct. Since $2\Delta_i = \varphi_2^* \hat{\ell}_i$, Δ_i is a component of the ramification divisor of φ_2 for every $i \in \{1, \dots, 5\}$.

Let R be the ramification divisor of $\varphi_2 : S \rightarrow S_2$. Denote by R_i the reduced and irreducible components of R such that $B_i = \varphi_2(R_i)$ is a curve, and let E_j be the components of R which are contracted by φ_2 . Then (cf. [2, Lemme 3.2])

$$K_S = \varphi_2^* K_{S_2} + \sum (r_i - 1) R_i + \sum e_j E_j,$$

where $e_j \geq 0$ is an integer and $r_i \geq 2$ is the ramification index of φ_2 at a generic point of R_i , i.e. the coefficient of R_i in $\varphi_2^* B_i$. Since K_S is ample we have $\sum E_j = 0$.

Whence, since $2K_S \equiv \varphi_2^*(-K_{S_2})$, we get

$$3K_S \equiv \sum (r_i - 1)R_i.$$

Let $e \subset S_2$ be the the image of the exceptional curve of μ_x , and let $E = \varphi^*e$. Then $E^2 = -4$ and $E \cdot K_S = 2$. By [16] there are no $(-n)$ -curves on S . Hence, since K_S is ample, $E = A_1 + A_2$, where A_1, A_2 are irreducible curves with $A_i \cdot K_S = 1$ and, by adjunction and the algebraic index theorem, $A_i^2 = -1$, $i = 1, 2$. Then $(A_1 + A_2)^2 = -4$ yields $A_1 \cdot A_2 = -1$, i.e. $A_1 = A_2$. In particular, A_1 is a component of R with ramification index 2.

It follows that the divisor

$$3K_S - A_1 - \sum_{i=1}^5 \Delta_i$$

is effective and A_1 does not belong to it. But then, since $4\Delta_i \cdot A_1 = \hat{L} \cdot E = 4(\hat{\ell} \cdot e) = 4$ for every i , we get

$$(3K_S - A_1 - \sum \Delta_i) \cdot A_1 = 3 + 1 - 5 < 0,$$

a contradiction. Therefore, $F \cdot M \neq 0$.

Suppose $M \cdot F = 1$. Thus M is isomorphic to $\mathbb{B}$, and hence $g(\mathbb{B}) = 3$. But then, by (1), we get $g(F) = 2$ which contradicts (2).

Therefore we are left with $M \cdot F = 2$. The restriction of $f : X \rightarrow \mathbb{B}$ to M is then a double cover $M \rightarrow \mathbb{B}$. Hence, since M is a smooth curve of genus 3 and $\mathbb{B}$ has genus 2, M is hyperelliptic [loc.cit].

Since $\hat{L}$ is isomorphic to M , $\varphi_{\hat{L}} : S \rightarrow \mathbb{P}^1$ is fibration with general fiber a hyperelliptic curve of genus 3. By [7] the bicanonical map of S factors through a double cover onto a rational surface. This completes the proof. $\square$

3. CONCLUSION

First of all we prove Theorem 0.1.

Proof of Theorem 0.1. Let d be the degree of φ_2 . Then $d = 2$ or 4 .

Suppose $d = 2$. Let σ be the involution induced by $\varphi_2 : S \rightarrow S_2$. Then φ_2 factors through σ and S_2 is birational to S/σ . Form Lemma 2.1 it follows that S_2 is rational.

Suppose that $d = 4$. Then, by Theorem 2.2, φ_2 factors through a double cover onto a rational surface. But this contradicts [18].

Therefore, $d = 2$, S_2 is a rational surface and, since K_S is ample, φ_2 is finite, i.e. a double cover. $\square$

The following theorem characterizes completely, from different points of view, minimal surfaces of general type with $p_g = q = 1$, $K_S^2 = 8$ and non bicanonical map.

Theorem 3.1. *Let S be a minimal surface of general type with $p_g = q = 1$ and $K_S^2 = 8$. The following are equivalent.*

- (1) *The bicanonical map is non birational.*
- (2) *The bicanonical map is a double cover onto a rational surface.*
- (3) *S is birational to a du Val double plane of type $\mathcal{D}_6$ (cf. [6]), i.e. the smooth minimal model of a double cover of $\mathbb{P}^2$ branched over the union B of a curve of degree 16 with 6 distinct lines through a point p , such that the essential singularities of B are the following:*
 - *p is a singular point of multiplicity 14,*
 - *there is a $[5, 5]$ -point on each line.*
- (4) *S is isomorphic to the quotient of a product $F \times C$ of two smooth curves of genus $g(F)$ and $g(C) = 3$ by a group G , where either*
 - *$g(F) = 3$ and $G = \mathbb{Z}_2 \times \mathbb{Z}_2$, or*
 - *$g(F) = 4$ and $G = S_3$, or*
 - *$g(F) = 5$ and $G = D_4$.*
- (5) *There exists a fibration $S \rightarrow \mathbb{P}^1$ whose general fiber is a smooth hyperelliptic curve of genus 3.*

Proof. Implication (1) $\Rightarrow$ (2) follows from Theorem 0.1, and (2) $\Rightarrow$ (1) is trivial. The implications (2) $\Rightarrow$ (3) and (3) $\Rightarrow$ (1) are proved in [6]. The equivalence (4) $\Leftrightarrow$ (2) follows from [18]. Finally, (1) $\Leftrightarrow$ (5) follows from [7]. $\square$

We remark that this classification is effective, that is these surfaces do exist. In fact, in [18] Polizzi proves that they form three components of the moduli space, one of dimension 5 and two of dimension 4. He also constructs examples as in (3), i.e. quotient of a product of curves.

REFERENCES

- [1] R. Accola *Topics in the theory of the Riemann surfaces*, Lecture Notes in Mathematics 1595 (1994),
- [2] A. Beauville, *L'application canonique pour les surfaces de type général*, Inventiones Math. **55** (1979), p.121-140.
- [3] A. Beauville, Appendice to *L'inégalité $p_g \geq 2q - 4$ pour les surfaces de type général*. Bull. Soc. Math. France **110** (1982), p. 343-346.
- [4] A. Beauville, *Complex algebraic surfaces*, Cambridge University Press 1996.
- [5] E. Bombieri, *Canonical models of surfaces of general type*, Inst. Hautes études Sci. Publ. Math., **42** (1973), p. 171-219.
- [6] G. Borrelli, *The classification of surfaces of general type with non birational bicanonical map*. J. Algebraic Geom. **16** (2007), 625-669.
- [7] G. Borrelli, *Bicanonical map of surfaces with $\chi = 1$ fibered by hyperelliptic curves of genus 3*. (submitted).
- [8] F.Catanese, *On a class of surfaces of general type*. in Algebraic Surfaces, CIME, Liguori (1991), 269-284.
- [9] C.Ciliberto, M. Mendes Lopes, *On surfaces with $p_g = q = 2$ and non-birational bicanonical map*, Adv. Geom. **2** (2002), no.3, 281-300.
- [10] I.Bauer, F. Catanese, R. Pignatelli, *Complex surfaces of general type: some recent progress*. in 'Global aspects of complex geometry' F.Catanese et al. editors, Springer Verlag.
- [11] F.Catanese, *Canonical rings and special surfaces of general type*. In: Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), 175-194. Proc. Sympos. Pure Math., **46**, Part 1. Amer. Math. Soc., Providence, RI (1987).
- [12] O.Debarre, *Inégalités numériques pour les surfaces de type général*, Bull. Soc. Math. France **110** (1982), 319-346.

- [13] M.Mendes Lopes, R.Pardini, *A survey on the bicanonical map of surfaces with $p_g = 0$ and $K \geq 2$* . In: Beltrametti, M.C. (ed) et al. Algebraic geometry. A volume in memory of Paolo Francia, 277-287. De Gruyter, Berlin (2002).
- [14] M.Mendes Lopes, R.Pardini, *The degree of the bicanonical map of a surface with $p_g = 0$* . Proc. Amer. Math. Soc. **135** (2007), 1279-1282.
- [15] Y. Miyaoka, *On the Chern numbers of surfaces of general type*. Invent. Math. **42**, (1977), 225-237.
- [16] Y. Miyaoka, *The maximum number of quotient singularities on surfaces with given numerical invariants*. Math. Ann. **268**, (1984), 159-171.
- [17] M. Nagata, *On rational surfaces I*. Mem. Coll. Sci. Univ. Kyoto **32**, 351-370 (1960).
- [18] F.Polizzi, *Surfaces of general type with $p_g = q = 1$, $K^2 = 8$ and bicanonical map of degree 2*. Trans. Amer. Math. Soc. **358**, (2006), 759-798.
- [19] I.Reider, *Vector bundles of rank 2 and linear systems on algebraic surfaces*. Ann. of Math., **127** (1988).
- [20] C. Rito, *On surface with $p_g = q = 1$ and non-ruled bicanonical involution*. Annali della Scuola Normale Superiore di Pisa. Classe di scienze, **6**, (2007), 81-102.
- [21] F. Serrano, *Isotrivial fibred surfaces*, Annali di Matematica pura e applicata vol. CLXXI (1996), 63-81.
- [22] G.Xiao, *Degree of the bicanonical map of a surface of general type*, Amer. J. of Math., **112** (5) (1990), 713-737.
- [23] G.Xiao, *Surfaces Fibrés en Courbes de Genre Deux*, Lecture Notes in Math. **1137**, (1985).
- [24] S. T. Yau, *On the Ricci curvature of a compact Kahler manifold and the complex Monge-Ampère equation*. I. Comm. Pure Appl. Math. **31**, no. 3, (1978), 339-411.

UNIVERSIDADE FEDERAL DE PERNAMBUCO,
 DEPARTAMENTO DE MATEMÁTICA,
 CIDADE UNIVERSITARIA, 50670-901 RECIFE (PE), BRASIL

E-mail address: borrelli@dmate.ufpe.br